

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
16 May 2002 (16.05.2002)

PCT

(10) International Publication Number
WO 02/38544 A2

(51) International Patent Classification⁷: C07D 209/00

(21) International Application Number: PCT/US01/50666

(22) International Filing Date: 26 October 2001 (26.10.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/247,304 10 November 2000 (10.11.2000) US
60/306,793 20 July 2001 (20.07.2001) US

(71) Applicant (for all designated States except US): ELI LILLY AND COMPANY [US/US]; Lilly Corporate Center, Indianapolis, IN 46285 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): JESUDASON, Cynthia, Darshini [LK/US]; 1090 Fleetwood Drive, Indianapolis, IN 46228 (US). SALL, Daniel, Jon [US/US]; 3601 Ostrom Court, Greenwood, IN 46143 (US). STEVENS, Freddie, Craig [US/US]; 139 Dahlia Lane, Indianapolis, IN 46217 (US). WERNER, John, Arnold [US/US]; 5486 Mark Court, Greenwood, IN 46142 (US).

(74) Agents: VOY, Gilbert, T. et al.; ELI LILLY AND COMPANY, Lilly Corporate Center, Indianapolis, IN 46285 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU (utility model), AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, CZ (utility model), DE, DE (utility model), DK, DK (utility model), DM, DZ, EC, EE, EE (utility model), ES, FI, FI (utility model), GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, I.K, I.R, I.S, I.T, I.U, I.V, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (utility model), SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

— as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

— as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: 3-SUBSTITUTED OXINDOLE BETA 3 AGONISTS

(57) Abstract: The present invention relates to a β_3 adrenergic receptor agonist of formula I: (I); or a pharmaceutical salt thereof; which is useful for treating Type II diabetes and/or obesity.

WO 02/38544 A2

- 1 -

3-SUBSTITUTED OXINDOLE β_3 AGONISTS

This application claims the benefit of U.S. Serial No.'s 60/247,304 and 60/306,793.

5

Field of Invention

The present invention is in the field of medicine, particularly in the treatment of Type II diabetes and obesity. More specifically, the present invention relates
10 to β_3 adrenergic receptor agonists useful in the treatment of Type II diabetes and obesity.

Background of the Invention

The current preferred treatment for Type II, non-
15 insulin dependent diabetes as well as obesity is diet and exercise, with a view toward weight reduction and improved insulin sensitivity. Patient compliance, however, is usually poor. The problem is compounded by the fact that there are currently no approved medications that adequately
20 treat either Type II diabetes or obesity.

One therapeutic opportunity that has recently been recognized involves the relationship between adrenergic receptor stimulation and anti-hyperglycemic effects. Compounds that act as β_3 receptor agonists have been shown
25 to exhibit a marked effect on lipolysis, thermogenesis and serum glucose levels in animal models of Type II (non-insulin dependent) diabetes.


The β_3 receptor, which is found in several types of human tissue including human fat tissue, has roughly 50%
30 homology to the β_1 and β_2 receptor subtypes yet is considerably less abundant. Stimulation of the β_1 and β_2

-2-

receptors can cause adverse effects such as tachycardia, arrhythmia, or tremors. An agonist that is selective for the β_3 receptor over the β_1 and β_2 receptors is, therefore, more desirable for treating Type II diabetes or obesity
 5 relative to a non-selective agonist.

However, recent studies have suggested the presence of an atypical beta receptor associated with atrial tachycardia in rats (*Br. J. of Pharmacol.*, 118:2085-2098, 1996). In other words, compounds that are not agonists of
 10 the β_1 and β_2 receptors can still modulate tachycardia through activation of a yet to be discovered β_4 or through some other unknown pathway.


A large number of publications have appeared in recent years reporting success in discovery of agents that
 15 stimulate the β_3 receptor. For example, U.S. Patent No. 5,786,356 discloses β_3 agonists of the formula:



(I)

wherein:


20 R_1 can be, among other things, a moiety of the formula:



A_1 and A_2 can be, among other things, NH, CH_2 , NCH_3 , or NCH_2CH_3 ; and

25 R_4 can be, among other things, a moiety of the formula:


-3-



Despite these recent developments, there remains a need to develop a selective β_3 receptor agonist which has minimal agonist activity against the β_1 and β_2 receptors and which displays a minimal propensity to cause atrial tachycardia.

Summary of Invention

The present invention relates to a compound of formula I:



I;

wherein:

R^1 is H, CN, halo, C_1 - C_6 alkyl, C_1 - C_4 haloalkyl, CO_2R^8 , $CONHR^8$, $NHCOR^8$, NHR^8 , OR^8 , SR^8 , SOR^8 , SO_2R^8 or SO_2NHR^8 ;

R^{1a} is H, halo or C_1 - C_6 alkyl;

R^2 is H, C_1 - C_6 alkyl or benzyl;

R^3 is C_1 - C_6 alkyl or benzyl;

or R^2 and R^3 combine with the carbon to which each are attached to form a C_3 - C_7 carbocyclic ring; provided that if R^3 is C_2 - C_6 alkyl or benzyl, then R^2 must be hydrogen;


R^4 is H or C_1 - C_6 alkyl;

R^5 and R^6 are independently H or C_1 - C_6 alkyl; or

-4-

R^5 and R^6 combine with the carbon to which each are attached to form a C_3 - C_6 carbocyclic ring;

or R^6 combines with X^1 , the carbon to which both are attached, and the phenyl group to which X^1 is attached to form a moiety selected from the group consisting of:



wherein:

m and n are independently 0, 1, 2, or 3

provided that the sum of $n + q$ is ≤ 5 and

that R^5 is H;

R^7 is hydrogen, optionally substituted phenyl or optionally substituted heterocycle;

R^8 is H or C_1 - C_6 alkyl;

X is OCH_2 , SCH_2 or a bond; and

X^1 is a bond or a C_1 - C_5 divalent hydrocarbon moiety; and

X^2 is O, S, NH, $NHSO_2$, SO_2NH , CH_2 or a bond; or a pharmaceutical salt thereof.

The present invention also relates to processes for preparing, as well as novel pharmaceutical formulations containing, a compound of formula I. In another embodiment, the pharmaceutical formulations of the present invention may be adapted for use in treating Type II diabetes and obesity and for agonizing the β_3 receptor.


The present invention also relates to methods for treating Type II diabetes and obesity, as well as a method

-5-

for agonizing the β_3 receptor employing a compound of formula I.

In addition, the present invention relates to a compound of formula I for use in treating Type II diabetes and obesity as well as a compound of formula I for use in agonizing the β_3 receptor. The present invention is further related to the use of a compound of formula I for the manufacture of a medicament for treating Type II diabetes and obesity as a well as for agonizing the β_3 receptor.

The present invention is also related to a compound of formula II:



II;

which is useful as an intermediate to prepare a compound of formula I.

Detailed Description

For the purposes of the present invention, as disclosed and claimed herein, the following terms are defined below.

The term "halo" represents fluoro, chloro, bromo, or iodo.

The term "C₁-C₆ alkyl" represents a straight, branched or cyclic hydrocarbon moiety having from one to six carbon atoms, e.g., methyl, ethyl, n-propyl, isopropyl, cyclopropyl, n-butyl, isobutyl, sec-butyl, t-butyl, cyclobutyl, pentyl, cyclopentyl, hexyl, cyclohexyl and the like. The term "C₁-C₄ alkyl" refers specifically to methyl, ethyl, n-propyl, isopropyl, cyclopropyl, n-butyl, isobutyl,

-6-

sec-butyl, t-butyl and cyclobutyl. A "C₁-C₄ haloalkyl" group is a C₁-C₄ alkyl moiety substituted with up to six halo atoms, preferably one to three halo atoms. An example of a haloalkyl group is trifluoromethyl. A "C₁-C₆ alkoxy" group is a C₁-C₆ alkyl moiety connected through an oxy linkage. A "C₁-C₄ alkoxy" group is a C₁-C₄ alkyl moiety connected through an oxy linkage.

The term "divalent hydrocarbon moiety" refers to a straight or branched chain of carbon atoms that may optionally have one or more points of unsaturation. Thus, a hydrocarbon diradical according to the present invention includes alkylene, alkenylene and alkylidene moieties. Examples include but are not intended to be limited to methylene, ethylene, propylene, butylene, -CH(CH₃)CH₂-, -CH(C₂H₅)CH₂-, -CH(CH₃)CH(CH₃)-, -CH₂C(CH₃)₂-, -CH₂CH(CH₃)CH₂-, -C(CH₃)₂CH₂-, -CH=CHCH₂-, -CH=CH-, -C≡CCH₂-, and the like.

The term "optionally substituted" as used herein means an optional substitution of one to three, preferably one or two groups independently selected from oxo, nitro, cyano, phenyl, benzyl, halo, C₁-C₆ alkyl, C₁-C₄ haloalkyl, COR⁹, NR¹⁰R¹⁰, NR¹⁰COR⁹, NR¹⁰SO₂R¹¹, OR¹⁰, OCOR⁹, OSO₂R¹¹, SR¹⁰, SOR¹¹, SO₂R¹¹ or SO₂NR¹⁰R¹⁰; wherein

R⁹ is H, C₁-C₆ alkyl, phenyl, benzyl, C₁-C₄ haloalkyl, NR¹⁰ar^{10a} or OR^{10a};

R¹⁰ and R^{10a} are independently H, C₁-C₆ alkyl or phenyl; or when two R¹⁰ or R^{10a} groups are attached to the same nitrogen atom, said R¹⁰ or R^{10a} groups, together with the nitrogen to which they are attached, may combine to form a piperidine, pyrrolidine, hexamethyleneimine or morpholine ring; and

-7-

R¹¹ is C₁-C₆ alkyl or phenyl.

The term "heterocycle" represents a stable, saturated, partially unsaturated, fully unsaturated or aromatic 5 or 6 membered ring, said ring having from one to
5 four heteroatoms that are independently selected from the group consisting of sulfur, oxygen, and nitrogen. The heterocycle may be attached at any point which affords a stable structure. Representative heterocycles include 1,3-dioxolane, 4,5-dihydro-1H-imidazole, 4,5-dihydrooxazole,
10 furan, imidazole, imidazolidine, isothiazole, isoxazole, morpholine, oxadiazole, oxazole, oxazolidinedione, oxazolidone, piperazine, piperidine, pyrazine, pyrazole, pyrazoline, pyridazine, pyridine, pyrimidine, pyrrole, pyrrolidine, tetrazole, thiadiazole, thiazole, thiophene and
15 triazole.

The term "suitable solvent" refers to any solvent, or mixture of solvents, inert to the ongoing reaction that sufficiently solubilizes the reactants to afford a medium within which to effect the desired reaction.

20 The term "patient" includes human and non-human animals such as companion animals (dogs and cats and the like) and livestock animals. Livestock animals are animals raised for food production. Ruminants or "cud-chewing" animals such as cows, bulls, heifers, steers, sheep,
25 buffalo, bison, goats and antelopes are examples of livestock. Other examples of livestock include pigs and avians (poultry) such as chickens, ducks, turkeys and geese. Yet other examples of livestock include fish, shellfish and crustaceans raised in aquaculture. Also included are exotic
30 animals used in food production such as alligators, water buffalo and ratites (e.g., emu, rheas or ostriches).
The preferred patient of treatment is a human.

The terms "treating" and "treat", as used herein, include their generally accepted meanings, i.e., preventing,

-8-

prohibiting, restraining, alleviating, ameliorating, slowing, stopping, or reversing the progression or severity of a pathological condition, or sequela thereof, described herein.

5 The terms "preventing", "prevention of", "prophylaxis", "prophylactic" and "prevent" are used herein interchangeably and refer to reducing the likelihood that the recipient of a compound of formula I will incur or develop any of the pathological conditions, or sequela
10 thereof, described herein.

 As used herein, the term "effective amount" means an amount of a compound of formula I that is capable of treating conditions, or detrimental effects thereof, described herein or that is capable of agonizing the β_3
15 receptor.

 The term "selective β_3 receptor agonist" means a compound that displays preferential agonism of the β_3 receptor over agonism of the β_1 or β_2 receptor. Thus, β_3 selective compounds behave as agonists for the β_3 receptor
20 at lower concentrations than that required for similar agonism at the β_1 and β_2 receptors. A β_3 selective compound also includes compounds that behave as agonists for the β_3 receptor and as antagonists for the β_1 and β_2 receptors.

 The term "pharmaceutical" when used herein as an
25 adjective means substantially non-deleterious to the recipient patient.

 The term "formulation", as in pharmaceutical formulation, is intended to encompass a product comprising the active ingredient(s) (compound of formula I), and the
30 inert ingredient(s) that make up the carrier, as well as any product which results, directly or indirectly, from combination, complexation or aggregation of any two or more

-9-

of the ingredients, or from dissociation of one or more of the ingredients, or from other types of reactions or interactions of one or more of the ingredients.

Accordingly, the pharmaceutical formulations of the present invention encompass any composition made by admixing a compound of the present invention and a pharmaceutical carrier.

The term "unit dosage form" refers to physically discrete units suitable as unitary dosages for human subjects and other non-human animals, each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect, in association with a suitable pharmaceutical carrier.

Because certain compounds of the invention contain an acidic moiety (e.g., carboxy), the compound of formula I may exist as a pharmaceutical base addition salt thereof. Such salts include those derived from inorganic bases such as ammonium and alkali and alkaline earth metal hydroxides, carbonates, bicarbonates, and the like, as well as salts derived from basic organic amines such as aliphatic and aromatic amines, aliphatic diamines, hydroxy alkamines, and the like.

Because certain compounds of the invention contain a basic moiety (e.g., amino), the compound of formula I can also exist as a pharmaceutical acid addition salt. Such salts include the salicylate, sulfate, pyrosulfate, bisulfate, sulfite, bisulfite, phosphate, mono-hydrogenphosphate, dihydrogenphosphate, metaphosphate, pyrophosphate, chloride, bromide, iodide, acetate, propionate, decanoate, caprylate, acrylate, formate, isobutyrate, heptanoate, propiolate, oxalate, malonate, succinate, suberate, sebacate, fumarate, maleate, 2-butyne-1,4 dioate, 3-hexyne-2, 5-dioate, benzoate, chlorobenzoate, hydroxybenzoate, methoxybenzoate, phthalate,

-10-

xylenesulfonate, phenylacetate, phenylpropionate,
phenylbutyrate, citrate, lactate, hippurate, β -
hydroxybutyrate, glycolate, maleate, tartrate,
methanesulfonate, propanesulfonate, naphthalene-1-sulfonate,
5 naphthalene-2-sulfonate, mandelate and like salts.
Preferred acid addition salts include the hydrochloride and
glycolate salts.

It is recognized that various stereoisomeric forms
of a compound of formula I exist. The compounds may be
10 prepared as racemates and can be conveniently used as such.
Therefore, the racemates, individual enantiomers,
diastereomers, or mixtures thereof form part of the present
invention. Unless otherwise specified, whenever a compound
is described or referenced in this specification all the
15 racemates, individual enantiomers, diastereomers, or
mixtures thereof are included in said reference or
description.

It is also recognized that various tautomeric
forms of a compound of formula I may exist, and all
20 tautomeric forms are part of the present invention. Unless
otherwise specified, whenever a compound is described or
referenced in this specification all tautomeric forms, or
mixtures thereof, are included in said reference or
description.

25

Preferred Compounds of the Invention

Certain compounds of the invention are
particularly interesting and are preferred. The following
listing sets out several groups of preferred compounds. It
30 will be understood that each of the listings may be combined
with other listings to create additional groups of preferred
compounds.

-11-

- a) R^1 is H, CN, halo, C_1 - C_4 alkyl, C_1 - C_4 haloalkyl, CO_2R^8 , $CONHR^8$, $NHCOR^8$, NHR^8 , OR^8 , SR^8 , SOR^8 , SO_2R^8 or SO_2NHR^8 ;
- b) R^1 is H, methyl, ethyl, CF_3 , chloro or fluoro;
- 5 c) R^1 is H, methyl, chloro or fluoro;
- d) R^1 is H or fluoro;
- e) R^1 is H;
- f) R^{1a} is H, halo or C_1 - C_4 alkyl;
- g) R^{1a} is H, methyl, ethyl, chloro or fluoro;
- 10 h) R^{1a} is H, methyl, chloro or fluoro;
- i) R^{1a} is H;
- j) R^2 is H and R^3 is methyl;
- k) R^2 and R^3 are both methyl;
- l) R^2 and R^3 combine with the carbon to which each are
15 attached to form a pentacyclic or hexacyclic ring;
- m) R^2 and R^3 combine with the carbon to which each are
attached to form a propacyclic or butacyclic ring;
- n) R^2 and R^3 are both methyl or combine with the carbon
to which each are attached to form a pentacyclic
20 ring;
- o) R^2 and R^3 combine with the carbon to which each are
attached to form a pentacyclic ring;
- p) R^4 is H or C_1 - C_4 alkyl;
- q) R^4 is H;
- 25 r) R^5 is H or C_1 - C_4 alkyl;
- s) R^6 is H or C_1 - C_4 alkyl;
- t) R^5 is H or methyl;
- u) R^6 is H or methyl;
- v) R^5 and R^6 are both methyl;

-12-

- w) R^7 is hydrogen, phenyl or heterocycle wherein said phenyl or heterocycle is optionally substituted one to three times independently with hydroxy, oxo, nitro, phenyl, benzyl, C_1 - C_4 alkoxy, COR^8 , $NHCO(C_1$ - C_4 alkyl), $NHCO(phenyl)$, $NHCO(benzyl)$, $OCO(C_1$ - C_4 alkyl), OCO_2R^8 and $OCONR^8R^8$;
- x) R^7 is selected from hydrogen, optionally substituted phenyl, pyridyl, thienyl and furanyl;
- y) R^7 is phenyl, pyridyl, thienyl or furanyl wherein said R^7 moieties are substituted one to three times with fluoro, chloro, cyano, hydroxy, methyl, ethyl, trifluoromethyl, methoxy, ethoxy, amino, CO_2CH_3 , $CO_2CH_2CH_3$, $CONR^8R^8$, SCH_3 , SCH_2CH_3 , $SOCH_3$, $SOCH_2CH_3$, SO_2CH_3 or $SO_2CH_2CH_3$;
- z) R^7 is phenyl, pyridyl, thienyl or furanyl wherein said R^7 moieties are substituted one to three times with fluoro, cyano, hydroxy, methyl, ethyl, methoxy, ethoxy, amino, carboxymethyl, carboxyethyl, $CONH_2$, CO_2CH_3 , $CO_2CH_2CH_3$, $CONH_2$, SCH_3 , SCH_2CH_3 , $SOCH_3$, $SOCH_2CH_3$, SO_2CH_3 or $SO_2CH_2CH_3$;
- aa) R^7 is phenyl, pyridyl, thienyl or furanyl wherein said R^7 moieties are substituted one to three times with fluoro, amino, CO_2CH_3 , $CO_2CH_2CH_3$, cyano, $CONH_2$, SO_2CH_3 or $SO_2CH_2CH_3$;
- bb) R^7 is phenyl, pyridyl, pyridazinyl or pyrimidinyl wherein said R^7 moieties are optionally substituted once or twice with chloro, cyano, $CONH_2$ or CO_2CH_3 ;
- cc) R^7 is phenyl, pyridyl, thienyl or furanyl wherein said R^7 moieties are substituted once with cyano or $CONH_2$;


-13-

- dd) R^7 is phenyl or pyridyl wherein said R^7 moieties are substituted once with cyano or CONH_2 ;
- ee) R^7 is pyridyl substituted once with cyano or CONH_2 ;
- ff) R^7 is 5-cyano or 5-carboxamido-pyrid-2-yl;
- 5 gg) R^7 is 4-cyano or 4-carboxamido-phenyl;
- hh) R^7 is 3-cyano or 3-carboxamido-pyrid-2-yl;
- ii) R^7 is 2-cyano or 2-carboxamido-phenyl;
- jj) R^8 is H or $\text{C}_1\text{-C}_4$ alkyl;
- kk) R^8 is hydrogen;
- 10 ll) X is connected to the indole ring system at the 4-position of said system;
- mm) X is OCH_2 ;
- nn) X^1 is a bond, methylene or ethylene;
- oo) X^1 is methylene;
- 15 pp) X^2 is at the para-position relative to X^1 ;
- qq) X^2 is a bond or O;
- rr) X^2 is O;
- ss) the compound of formula I is an acid addition salt;
- tt) the compound of formula I is the hydrochloride salt;
- 20 uu) the compound of formula I is the glycolate salt.

Synthesis

The compound of formula I may be prepared as described in the following Schemes and Examples.

-14-

Scheme 1

The reaction of Scheme 1 may be carried out under
 5 conditions appreciated in the art for the amination of
 epoxides. For example, the epoxide of formula II may be
 combined with an amine of formula III in a lower alcohol,
 dimethylformamide, dimethylsulfoxide, or acetone, preferably
 ethanol, isopropanol, n-butanol or t-butanol, at room
 10 temperature to the reflux temperature of the reaction
 mixture, preferably between 40°C - 90°C. The reaction may
 also be carried out under conditions generally described in
 Atkins, *et al.*, *Tet. Let.*, 27:2451, 1986. These conditions
 include mixing the reagents in the presence of
 15 trimethylsilyl acetamide in a polar aprotic solvent such as
 acetonitrile, dimethylformamide, acetone, dimethylsulfoxide,
 dioxane, diethylene glycol dimethyl ether, tetrahydrofuran,
 or other polar aprotic solvents in which the reagents are
 soluble.


20 The epoxide starting materials employed in Scheme
 1 may be prepared by techniques recognized and appreciated
 by one skilled in the art. See, *e.g.*, references cited
 below in the Preparations section for representative and/or
 analogous procedures for preparing the epoxides of formula

-15-


II. To illustrate, epoxides of formula II, where X is OCH₂ or SCH₂ and where R² and R³ combine with the carbon to which each are attached to form a C₃-C₇ carbocyclic ring, may be prepared according to the procedure detailed in Scheme 2

5 wherein R¹² is chloro, bromo or iodo, X' is O or S, and X'' is OCH₂ or SCH₂ and p is 0-4.

In addition, epoxides of formula II, where X is OCH₂ or SCH₂ and where R² and R³ do not combine with the carbon to which each are attached to form a carbocyclic ring, may be prepared according to the procedure detailed in
 10 Scheme 3 below where R^{2'} is C₁-C₆ alkyl or benzyl.

Scheme 2

-16-

Scheme 3


5 The compounds of formula II(a) and II(b) may be prepared by reacting equimolar amounts of a compound of formula VIII or VIII(a), respectively, with (2S)-(+)-glycidyl 3-nitrobenzenesulfonate. The reaction is typically performed in an inert solvent such as acetone and in the presence of a slight excess of a weak base, such as

 10 potassium carbonate. The suspension may then be heated at reflux for 16-20 hours with stirring to provide a compound of formula II(a) or II(b).

15 The cyclopropyl derivatives are prepared by a slightly modified procedure which involves treating the O- and N-diacetylated oxindole with 1,2 dibromoethane and potassium carbonate in dimethylsulfoxide.

20 The amino starting materials employed in Scheme 1 (formula III compound) may also be prepared by techniques recognized and appreciated by one skilled in the art. For example, an amine of formula III, where X² is O, may be prepared according to the procedure detailed in Scheme 4.

-17-

Scheme 4

A compound of formula XII may be prepared by reacting an arylalkyl alcohol of formula X with excess (5 mol/equivalent) formula X compound by methods well known in the art (see, e.g., *Sh. Prikl. Kin.*, 45:1573-77, 1972). The reaction may also be carried out by mixing the reagents in an aprotic solvent, preferably diglyme, and adding potassium t-butoxide (0.5 mol/equivalent). The reaction is typically heated at reflux until water present in the reaction mixture is removed (generally 2-8 hours). A compound of formula XIII may then be prepared by hydrogenation of the corresponding compound of formula XII over a precious metal catalyst. The hydrogenation can be affected at between 20 and 60 psi of hydrogen (preferably 50 psi), and with a variety of solvents (preferably methanol/acetic acid), temperatures (preferably 50°C), and catalysts (preferably 5% palladium on carbon wetted with ethanol denatured with toluene) well known in the art.

A skilled artisan will appreciate that a compound of formula XIII could be coupled with a wide variety of aryl halides to yield the claimed ethers. The coupling can be carried out according to procedures well known in the art

-18-

and is preferably performed by mixing the starting materials in N,N-dimethylacetamide and toluene in the presence of potassium carbonate. The reaction is typically then heated to reflux for 5 to 24 hours to effect the reaction and to
 5 remove water present in the reaction mixture.

Compounds of formula $R^{2'}-I$, R^3-I , IV, X, XI and $I-(CH_2)_p-I$, are either commercially available, known in the art, or can be prepared by methods known in the art or described herein.

10 The following Preparations, Examples and Formulations are provided so that the invention might be more fully understood. They should not be construed as limiting the invention in any way.

15 Preparations

Epoxides of Formula II

Epoxides 1-9 are prepared for use as described in Scheme 1. These epoxides are pictured below in Table 1.

20

Table 1










 <p>1</p>	 <p>2</p>	 <p>3</p>
--	--	--

Table 1 (continued)

 <p>4</p>	 <p>5</p>	 <p>6</p>
 <p>7</p>	 <p>8</p>	 <p>9</p>

Epoxide 1

5

Hydrazine monohydrate (265 mL, 5.46 mol) is added to a flask containing 4-methoxyisatin (484.26 g, 2.73 mol), sodium acetate (22.31 g, 0.27 mol) and dimethylformamide (2.4 L). The temperature rose from 19.2°C to 31.9°C within 10 5 minutes, and when the exotherm had ceased the solution is further heated to 110°C over 40 minutes. Vigorous off-gassing is observed, and the temperature rose above the set point slowly. After 30 minutes the temperature stabilized at 110°C and the gas evolution tapered off. The solution is 15 kept at 110°C for 40.5 hours at which point the heating mantle is removed and the solution is allowed to cool. At 40°C, the solution is seeded with 100 mg of 4-methoxyindole, and the contents are cooled to 17°C with an ice bath. The resulting mixture is poured into water (6.2 L) which had 20 been pre-cooled to 6°C. The flask is rinsed with water (600 mL), and the thick slurry is cooled to 4°C before being filtered through polypropylene. The filter cake is rinsed

-20-

with water (3 x 1.5 L) and the solid is dried in a 50°C vacuum oven to provide 353 g of 4-methoxyindolin-2-one.

A mixture of tetrahydrofuran (990 mL), 4-methoxyindolin-2-one (*J. Chem. Soc.*, 3904, 1952; *Tetrahedron*, 24:6093, 1968, 39.60 g, 242.7 mmol), and N,N,N',N'-tetramethylethylenediamine (56.40 g, 485.4 mmol) is cooled to -78°C. *N*-butyllithium (379.2 mL, 606.7 mmol, 1.6M in hexanes) is added over 1 hour so as to maintain the reaction temperature between -70°C and -69°C. After stirring the mixture for 50 minutes at -78°C, iodomethane (36.17 g, 254.8 mmol) is added over 50 minutes at -78°C to -64°C. The reaction mixture is allowed to warm to -40°C and iodomethane (48.23 g, 0.3398 mol) is added incrementally over 1 hour until the starting material is less than 3 percent by HPLC (retention 7.3 minutes; Zorbax SB-Phenyl (4.6 mm x 25 cm); solvent A - trifluoroacetic acid solution (0.1%, v/v); solvent B - acetonitrile; 1.0 mL/min; 254 nm; gradient method: 0.00 - 2.00 minutes, 30% A, 70% B; 2.00 - 7.00 minutes, linear ramp from 30% A, 70% B to 80% A, 20% B; 7.00 - 15.00 minutes, 80% A, 20% B). Methanol (20 mL) is added to the reaction mixture over 15 minutes at -60°C to -50°C. Aqueous hydrochloric acid (1N, 250 mL) is added allowing the temperature to rise to -4°C. The solution is transferred to a separatory funnel with ethyl acetate (500 mL), aqueous hydrochloric acid (1N, 250 mL) and a saturated sodium chloride solution (250 mL). After separating, the organic layer is extracted twice with aqueous hydrochloric acid (1N, 250 mL) and the aqueous layer is extracted with ethyl acetate (250 mL). The combined organic layers are washed with a saturated sodium chloride solution (250 mL) then dried over magnesium sulfate. After filtration, the solvents are removed by rotary evaporation to give 43.58 g (91.6%) of 4-methoxy 3-methyl-1,3-dihydro-

-21-

indol-2-one as a solid, mp 128-130 °C. MS (ES+) m/z 178 ($M^+ + 1$).

4-Methoxy-3-methyl-1,3-dihydro-indol-2-one is demethylated according to Representative Procedure 1 below to form 4-hydroxy-3-methyl-1,3-dihydro-indol-2-one. 4-Hydroxy-3-methyl-1,3-dihydro-indol-2-one is coupled to (2S)-glycidyl 3-nitrobenzenesulfonate to form the title epoxide according to Representative Procedure 2 below.

10

Epoxide 2

A solution of 4-methoxy-3-methyl-1,3-dihydro-indol-2-one (41.73 g, 235.5 mmol) and tetrahydrofuran (1000 mL) is cooled to -78°C. Potassium bis(trimethylsilyl)amide (989.0 mL, 494.3 mmol, 0.5 M in toluene) is added over 45 minutes so as to maintain the reaction temperature between -71°C and -66°C. Iodomethane (36.71 g, 258.7 mmol) is added over 15 minutes between -75°C and -70°C. The mixture is stirred at -78°C for 1 hour then -60°C for 30 minutes. Methanol (25 mL) is added at -60°C. Aqueous hydrochloric acid (1N, 420 mL) is added rapidly allowing the temperature to rise to -7°C. The solution is transferred to a separatory funnel with aqueous hydrochloric acid (1N, 420 mL) and toluene (50 mL). The organic layer is extracted with a saturated sodium bicarbonate solution (250 mL), washed with a saturated sodium chloride solution (200 mL), dried over magnesium sulfate, and filtered. The filtrate is stirred with DARCO (25 g) for 1 hour then filtered. The filtrate is concentrated by rotary evaporation. The residue is refluxed for 1 hour in *tert*-butyl methyl ether (300 mL). After distilling out 100 mL of solvent, the slurry is stirred for 10 hours at 24°C. The solid is isolated by vacuum filtration rinsing twice with cold

-22-

(-40°C) *tert*-butyl methyl ether. After vacuum drying for 12 hours at 50°C/5 Torr, 23.9 g (58%) of 4-methoxy-3,3-dimethyl-1,3-dihydro-indol-2-one is obtained. mp 143-144 °C. MS (ES+) m/z 192 ($M^+ + 1$).

- 5 Pyridine hydrochloride (51.10 g, 442.2 mmol) and 4-methoxy-3,3-dimethyl-1,3-dihydro-indol-2-one (21.61 g, 113.0 mmol) are combined and stirred in a melt at 220°C for 45 minutes. Heating is removed and when the mixture cooled to 100°C, water (60 mL) is added, followed by ethyl acetate
10 (150 mL) at 65°C. The layers are separated and the aqueous phase is extracted five times with ethyl acetate (50 mL). The combined organic layers are extracted first with a combination of aqueous hydrochloric acid (1N, 10 mL) and a saturated solution of sodium chloride (10 mL), then with a
15 combination of a saturated solution of sodium bicarbonate (10 mL) and a saturated solution of sodium chloride (10 mL). After drying over magnesium sulfate, the solvent is removed via rotary evaporation. The crude material is dissolved in ethyl acetate (200 mL) and hexane (200 mL) at 60°C.
20 Crystallization is allowed to begin slowly over 1 hour at 50°C. After stirring for 12 hours at 24°C, the slurry is cooled to 0°C for 1 hour, then vacuum filtered washing twice with cold 1:1, ethyl acetate:hexane (35 mL, 0°C) to provide 13.8 g (98.6% purity, 68% yield) of 4-hydroxy-3,3-dimethyl-
25 1,3-dihydro-indol-2-one, mp 224-225 °C. MS (ES+) m/z 178 ($M^+ + 1$). From the filtrate a second crop (5.0 g, 98.4% purity, 25% yield) is obtained to give a total yield of 93%.

- Acetone (250 mL), 4-hydroxy-3,3-dimethyl-1,3-dihydro-indol-2-one (17.56 g, 99.1 mmol), powdered potassium
30 carbonate (28.70 g, 208.1 mmol) and (2S)-(+)-glycidyl 3-nitrobenzenesulfonate (26.20 g, 101.1 mmol) are combined and the resulting mixture is refluxed for 22.5 hours. The solvent is removed by rotary evaporation and the crude material is purified by flash chromatography (500 g SiO₂;

-23-

1:1, ethyl acetate:hexane). During solvent removal from the product fractions, hexane (200 mL) is added and the product is crystallized in two crops. Isolation by vacuum filtration followed by vacuum drying at 50°C/5 Torr for 12 hours provided 11.0 g (47%) of the title epoxide, mp 156-158°C. MS (ES+) m/z 234 ($M^+ + 1$).

Epoxide 3

10 A mixture of 4-methoxyoxindole (3.0 g, 18.4 mmol) and piperidine (3.5 mL, 36.89 mmol) in acetone (100 mL) is heated at reflux for 18 hours. The mixture is concentrated to dryness and triturated with methanol (20 mL). Filtration removed 800 mg of unreacted 4-methoxyoxindole. The filtrate
15 is preadsorbed on silica gel (5 g), and purified by flash chromatography (120 g SiO₂, 25% ethyl acetate:hexane to 50% ethyl acetate:hexane) to provide 700 mg (19%) of 3-isopropylidene-4-methoxy-1,3-dihydro-indol-2-one. MS (ESI+) 204.

20 A mixture of 3-isopropylidene-4-methoxy-1,3-dihydro-indol-2-one (700 mg, 3.4 mmol) and 5% palladium on carbon (88 mg) in ethanol (25 mL) is hydrogenated at 60 psi overnight. Filtration and removal of the solvent *in vacuo* provided 560 mg (79%) of 3-isopropyl-4-methoxy-1,3-dihydro-indol-2-one. MS (ESI+) 206.

25 3-Isopropyl-4-methoxy-1,3-dihydro-indol-2-one is demethylated according to Representative Procedure 1 below in 45% yield. MS (ESI+) 192. 3-Isopropyl-4-hydroxy-1,3-dihydro-indol-2-one is coupled to (2S)-glycidyl 3-
30 nitrobenzenesulfonate to form the title epoxide according to Representative Procedure 2 below.

-24-

Epoxide 4

3-Isobutyl-4-methoxy-1,3-dihydro-indol-2-one is prepared from 4-methoxyoxindole (3.1 mmol) and 1-iodo-2-methylpropane (3.1 mmol) substantially as described above for Epoxide 1 and Epoxide 2. The compound is purified by flash chromatography (SiO₂, 25% ethyl acetate:hexane) to obtain 180 mg (27%) of the desired compound. MS (ESI+) 220.

3-Isobutyl-4-methoxy-1,3-dihydro-indol-2-one is demethylated according to Representative Procedure 1 below in 80% yield. MS (ESI+) 206. 3-Isobutyl-4-hydroxy-1,3-dihydro-indol-2-one is coupled to (2S)-glycidyl 3-nitrobenzenesulfonate to form the title epoxide according to Representative Procedure 2 below. MS (ESI+) 262.

Epoxide 5

3-Benzyl-4-methoxy-1,3-dihydro-indol-2-one is prepared from 4-methoxyoxindole (4.8 mmol) and benzyl bromide (5.3 mmol) substantially as described above for Epoxide 4. The compound is purified by flash chromatography (SiO₂, 25% ethyl acetate:hexane) to obtain the desired compound (36%). MS (ESI+) 254.

3-Benzyl-4-methoxy-1,3-dihydro-indol-2-one is demethylated according to Representative Procedure 1 below in 73% yield. MS (ESI+) 240. 3-Benzyl-4-hydroxy-1,3-dihydro-indol-2-one is coupled to (2S)-glycidyl 3-nitrobenzenesulfonate to form the title epoxide according to Representative Procedure 2 below. MS (ESI+) 296.

Epoxide 6

To a mixture of 4-(acetyloxy)-1,3-dihydro-2H-indol-2-one (*J. Med. Chem.*, 38:2802-2808, 1995; 1.44 g, 7.5

-25-

mmol) and sodium carbonate (5.3 g, 50 mmol) in tetrahydrofuran (50 mL) is added acetic anhydride (4.3 mL, 45 mmol). The mixture is stirred at ambient temperature for 18 hours, and partitioned between ethyl acetate and water. 5 After drying (magnesium sulfate), the solvent is removed in vacuo. The crude material is preadsorbed on silica gel (10 g), and purified by flash chromatography (90 g SiO₂, 10% ethyl acetate:hexane to 70% ethyl acetate:hexane, linear gradient over 30 minutes) to provide 480 mg (27%) of acetic 10 acid 1-acetyl-2-oxo-2,3-dihydro-1H-indol-4-yl ester as a white solid.

By a procedure substantially similar to that described in *Heterocyclic Chem.*, 31:1513, 1994, a mixture of acetic acid 1-acetyl-2-oxo-2,3-dihydro-1H-indol-4-yl ester 15 (0.48 g, 2.1 mmol) and potassium carbonate (0.57 g, 4.1 mmol) in dimethylsulfoxide (20 mL) is treated with 1,2-dibromoethane (0.39 mL, 4.5 mmol). The mixture is stirred at room temperature for 18 hours, and partitioned between diethylether and water. The aqueous phase is extracted with 20 diethylether (2x), and the combined organic phase is washed with brine, dried (magnesium sulfate), filtered and concentrated to dryness. The crude material is preadsorbed on silica gel (7 g), and purified by flash chromatography (90 g SiO₂, 10% ethyl acetate:hexane to 70% ethyl 25 acetate:hexane, linear gradient over 40 minutes) to yield 420 mg (79%) of 3-spirocyclopropane-4-methoxy-1,3-dihydro-indol-2-one. MS (ESI+) 260.

A mixture of 3-spirocyclopropane-4-methoxy-1,3-dihydro-indol-2-one (0.42 g, 1.6 mmol) in aqueous sulfuric acid (3N, 15 mL) and tetrahydrofuran (15 mL) is heated at 30 reflux for 2 hours. The mixture is partitioned between ethyl acetate and water. The aqueous phase is extracted twice with ethyl acetate, and the combined organic phase is washed with brine, and dried (magnesium sulfate). After

-26-

removal of the solvent under reduced pressure, the crude material is preadsorbed on silica gel (5 g), and purified by flash chromatography (40 g SiO₂, 10% ethyl acetate:hexane to 70% ethyl acetate:hexane, linear gradient over 40 minutes) to yield 140 mg (50%) of 3-spirocyclopropane-4-hydroxy-1,3-dihydro-indol-2-one. MS (ESI+) 176.

3-Spirocyclopropane-4-hydroxy-1,3-dihydro-indol-2-one is coupled to (2S)-glycidyl 3-nitrobenzenesulfonate to form the title epoxide according to Representative Procedure 2 below. MS (ESI+) 232.

Epoxide 7

A suspension of 4-methoxy-1,3-dihydro-indol-2-one (3.7 g, 22.9 mmol) in tetrahydrofuran (110 mL) is cooled to -78°C using a dry ice/acetone bath and treated with N,N,N',N'-tetramethylethylenediamine (8.6 mL, 57.1 mmol). N-butyllithium (36.0 mL, 57.1 mmol, 1.6 M in hexanes) is added dropwise, and the mixture stirred at this temperature for 30 minutes. 1,3-Diodopropane (13.2 mL, 114.5 mmol) is added and the mixture is slowly warmed to room temperature and stirred for 18 hours. The reaction is quenched by addition of methanol (50 mL), followed by 1N aqueous hydrochloric acid to dissolve the resulting solids. The resulting mixture is partitioned between ethyl acetate and brine. The aqueous phase is then extracted twice with ethyl acetate, and the combined organic phases are washed with brine, dried (magnesium sulfate), and concentrated. The crude material is preadsorbed on silica gel (15g) and purified by flash chromatography (90 g silica gel, 15% ethyl acetate/hexane to 50% ethyl acetate/hexane with a linear gradient over 45 minutes to afford 1.55 g of 3-spirocyclobutane-4-methoxy-1,3-dihydro-indol-2-one (33%). MS (ESI+) 204.

-27-

To a solution of 3-spirocyclobutane-4-methoxy-1,3-dihydro-indol-2-one (1.55 g, 7.6 mmol) in dichloromethane (100 mL) at -78°C is added dropwise a solution of boron tribromide in dichloromethane (1M, 38 mL, 38 mmol). After
5 the mixture is stirred at -78°C for 1 hour, the cooling bath is removed, and the mixture is allowed to warm to room temperature and stir overnight. The reaction is quenched by the addition of ice and water. The mixture is extracted three times with ethyl acetate, and the combined organic
10 phases are dried (magnesium sulfate), and concentrated. The crude material is preadsorbed on silica gel (7 g), and purified by flash chromatography (40 g silica gel, 25% ethyl acetate:hexane to 60% ethyl acetate:hexane, linear gradient over 45 minutes) to yield 750 mg (52%) of 3-
15 spirocyclobutane-4-hydroxy-1,3-dihydro-indol-2-one. MS (ESI+) 190.

3-Spirocyclobutane-4-hydroxy-1,3-dihydro-indol-2-one is coupled to (2S)-glycidyl 3-nitrobenzenesulfonate to form the title epoxide according to Representative Procedure
20 2 below. MS (ESI+) 246.

Epoxide 8

To a mixture of tetrahydrofuran (1000 mL) and 4-methoxy-1,3-dihydro-indol-2-one (48.30 g, 296.0 mmol) cooled
25 to -65°C was added *N,N,N',N'*-tetramethylethylenediamine (89.02 g, 766.0 mmol) maintaining the temperature between -65°C and -63°C. After cooling further to -75°C, *N*-butyllithium (478.8 mL, 766.0 mmol, 1.6 M in hexanes) is
30 added over 1 hour so as to maintain the temperature between -75°C and -73°C. After stirring the mixture for 30 minutes at -72°C, 1,4-dibromobutane (330.82 g, 1.5321 mol) is added over 1 hour between -72°C and -62°C. The solution is stirred at -33°C for 15 hours then at 24°C to 30°C for 5

-28-

hours. Methanol (63 mL) is added to the mixture and the pH is adjusted to 6 with concentrated hydrochloric acid (50 mL), 3N aqueous hydrochloric acid (50 mL) and 1N aqueous hydrochloric acid (500 mL). Ethyl acetate (1000 mL) and a saturated solution of sodium chloride (300 mL) are added. The organic layer is separated and extracted with a combination of a saturated solution of sodium bicarbonate (500 mL) plus a saturated solution of sodium chloride (300 mL) then dried over magnesium sulfate. After filtration the solvent is removed by rotary evaporation at 44°C/5 Torr until no further distillate is obtained. Heptane (500 mL) is added and removed twice by rotary evaporation. Heptane (250 mL) is added and the mixture is stirred at 24°C after which the solid is collected by vacuum filtration and rinsed three times with heptane (100 mL). After vacuum drying at 60°C/5 Torr for 14 hours, 53.1 g (94.3% purity, 78% yield) of 3-spirocyclopentane-4-methoxy-1,3-dihydro-indol-2-one is obtained as a solid, mp 168-169 °C. MS (ES+) m/z 218 ($M^+ + 1$). From the filtrate a second crop (3.6 g, 96.3% purity, 6% yield) is obtained to give a total yield of 84%.

Alternatively, 3-spirocyclopentane-4-methoxy-1,3-dihydro-indol-2-one can be prepared as follows. A slurry of 4-methoxyindolin-2-one (200 g, 1.2 mol) and tetrahydrofuran (2.6 L) is cooled to -70°C in a dry ice/acetone bath. A solution of sodium bis(trimethylsilyl)amide and tetrahydrofuran (4.9 L, 1M soln) pre-cooled to 4°C is added to this slurry over 1.5 hours while maintaining a temperature of ≤ -69 °C. The resulting solution is maintained at this temp for 20 minutes post addition at which point 1,4 dichlorobutane (347 g, 2.7 mol) is added in one portion. The cooling bath is removed and the solution is allowed to warm to 30°C over 4.5 hours. The solution is then heated to 35°C for an additional 15.5 hours. Quenching is effected with methanol (260 mL) followed by water (3 L),

-29-

and lastly, after installation of an ice bath, concentrated hydrochloric acid (640 mL) until the pH of the mixture is 2. The layers are separated and the aqueous layer is rinsed with ethyl acetate (1.6 L), and the combined organic layers
5 are washed with 1N aqueous hydrochloric acid/saturated sodium chloride (1.8 L/1.0 L). Again the aqueous layer is back-extracted with ethyl acetate (1.2 L). The combined organic layers are rinsed with saturated sodium
10 bicarbonate/saturated sodium chloride (1.2 L/1.2 L), dried (magnesium sulfate, 170 g) and filtered. The filtrate is then transferred to a rotary evaporator where a solvent exchange with heptane (4 L) is performed. When the slurry volume approaches 2 L it is cooled, filtered and dried in a 50°C vacuum oven overnight to provide a granular orange
15 solid (241.3, 91 %) which is used without further purification.

Pyridine hydrochloride (127.6 g, 1.1042 mol) and 3-spirocyclopentane-4-methoxy-1,3-dihydro-indol-2-one (48.29 g, 0.2223 mol) are combined and stirred in a melt at
20 220°C for 80 minutes. Heating is removed and at 100°C, water (150 mL) is added followed by ethyl acetate (300 mL) at 60°C. The layers are separated and the aqueous phase is extracted four times with ethyl acetate (100 mL). The combined organic layers are extracted first with a
25 combination of aqueous hydrochloric acid (1N, 100 mL) and a saturated solution of sodium chloride (100 mL), then with a combination of a saturated solution of sodium bicarbonate (100 mL), a saturated solution of sodium chloride (100 mL), and water (200 mL). After drying over magnesium sulfate,
30 the solvent is removed by rotary evaporation. The crude material is purified by flash chromatography (5 kg silica gel; 5.4% methanol, 43.3% heptane, 51.3% methylene chloride) to provide 38.7 g (85%) of 3-spirocyclopentane-4-hydroxy-

-30-

1,3-dihydro-indol-2-one as a solid, mp 215-216 °C. MS (ES+) m/z 204 ($M^+ + 1$).

Acetone (750 mL), 3-spirocyclopentane-4-hydroxy-1,3-dihydro-indol-2-one (37.76 g, 0.1858 mol), powdered
5 potassium carbonate (53.92 g, 0.3901 mol) and (2S)-(+)-glycidyl 3-nitrobenzenesulfonate (50.57 g, 0.1951 mol) are combined and the resulting yellow mixture refluxed for 20 hours. After the solvent is removed by rotary evaporation, the resulting solid is dissolved in ethyl acetate (750 mL),
10 water (2500 mL), and a saturated solution of sodium chloride (100 mL). The aqueous layer is separated and extracted with ethyl acetate (250 mL). The combined organic layers are extracted with a saturated solution of sodium chloride (200 mL), dried over sodium sulfate, and concentrated by rotary
15 evaporation. The crude material is purified by flash chromatography (5 kg silica gel; 12 L of 3.0% methanol, 55.0% heptane, 42.0% methylene chloride; then 9 L of 5.0% methanol, 47.5% heptane, 47.5% methylene chloride). After
vacuum drying at 50°C/5 Torr for 12 hours, 39.0 g (81%) of
20 the title epoxide is obtained as a solid, mp 153-154 °C. MS (ES+) m/z 260 ($M^+ + 1$).

Epoxide 9

25 A suspension of 4-methoxy-1,3-dihydro-indol-2-one (2.0 g, 212.3 mmol) in tetrahydrofuran (60 mL) is cooled to -78°C using a dry ice/acetone bath and treated with *N,N,N',N'*-tetramethylethylenediamine (3.7 mL, 24.6 mmol). *N*-butyllithium (15.4 mL, 24.6 mmol, 1.6 M in hexanes) is
30 added dropwise, and the mixture stirred at this temperature for 30 minutes. The mixture is warmed to -20°C, and 1,5-diiodopentane (9.2 mL, 61.5 mmol) is added, the mixture slowly warmed to room temperature and stirred for 18 hours. The reaction is quenched by addition of a saturated aqueous

-31-

ammonium chloride solution and partitioned between ethyl acetate and 1N aqueous hydrochloric acid. The aqueous phase is then extracted twice with ethyl acetate, and the combined organic phases are washed with brine and dried (magnesium sulfate). After removal of the solvent, the crude material is purified by flash chromatography (90 g silica gel, hexane to 50% ethyl acetate:hexane) to yield 1.13 g (40%) of 3-spirocyclohexane-4-methoxy-1,3-dihydro-indol-2-one. MS (ESI+) 232.

3-Spirocyclohexane-4-methoxy-1,3-dihydro-indol-2-one is demethylated according to Representative Procedure 1 below in 32% yield. MS (ESI+) 218. 3-Spirocyclohexane-4-hydroxy-1,3-dihydro-indol-2-one is coupled to (2S)-glycidyl 3-nitrobenzenesulfonate to form the title epoxide according to Representative Procedure 2 below.

Representative Procedure 1: Demethylation

An intimate mixture of the 4-methoxyindole and pyridine hydrochloride (approximately 10 g per gram oxindole) is heated for 15-45 minutes at 210-220°C in an oil bath. The reaction is cooled to room temperature, and the solid is added to a mixture of ice and ammonium hydroxide solution. This water mixture is evaporated, and the crude material preabsorbed on silica gel. A flash column in 25-50% ethyl acetate/hexanes provided the desired phenols in yields of 32-94%

Representative Procedure 2: Epoxide Formation

A mixture of the 4-hydroxyoxindole (1 equivalent), (2S)-glycidyl 3-nitrobenzenesulfonate (1.2 equivalents), potassium carbonate (1.2 equivalents) and acetone (resulting solution 0.2M in oxindole) is refluxed for 16 hours, cooled to room temperature and the solids removed via filtration.

-32-

The filtrate is concentrated, redissolved in ethyl acetate and extracted several times with water. The organic layer is concentrated and the crude product could be purified by flash chromatography (25% ethyl acetate/hexanes) to give the
 5 desired epoxides. However, in most cases the epoxide is used in the subsequent reactions without further purification.

Amines of Formula III

10 Amines 1-38 are prepared for use as described in Scheme 1. These amines are pictured below in Table 2.

Table 2








































			
R ⁷ =	R ⁷ =	R ⁷ =	R ⁷ =
 1	 2	 3	 4
 5	 6	 7	 8
 9	 10	 11	 12

Table 2 (continued)

 13	 14	 15	 16
 17	 18	 19	 20
 21	 22	 23	 24
 25	 26	 27	 28
 29	 30	 31	 32
 33	 34	 35	 36
	 37	 38	

-34-

Amines 1 and 9 may be prepared according to procedures detailed in U.S. Serial No. 09/068,192, the teachings of which are herein incorporated by reference. Amines 10 and 25 may be prepared by a procedure substantially similar to that described for Amine 1. Amines 2, 3 and 8 may be prepared according to procedures detailed in U.S. Patent No. 5,977,154, the teachings of which are herein incorporated by reference. Amines 26, 28 and 37 may be prepared by a procedure substantially similar to that described for Amine 8. Amine 32 may be prepared by a procedure substantially analogous to Amine 3. Amine 38 may be prepared according to procedures detailed in U.S. Patent No. 5,840,738, the teachings of which are herein incorporated by reference.

15

Amine 4

4-(2-Amino-2-methylpropyl)phenol (50.8 g, 225 mmol), 2-chloro-3-cyanopyridine (30.8 g, 222 mmol), potassium carbonate (77.7 g, 562 mmol, powdered), N,N-dimethylacetamide (609 ml), and isooctane (122 ml) are combined and heated to reflux. The water formed during the reaction is removed azeotropically via a Dean-Stark trap. After about 1-2 hours the reaction is complete. The slurry is cooled to 30°C and filtered. The filter cake is washed with N,N-dimethylacetamide (250 ml) and the combined organic fractions are concentrated by rotary evaporation at 80°C. The resulting dark green oil is dissolved in dichloromethane (580 ml), and washed with water (160 ml). The phases are separated and the organic phase washed with water (250 ml). Water (1 L) is added to the organic phase and the pH adjusted to 1 with 12N aqueous hydrochloric acid (about 25 ml). The phases are separated and the acidic aqueous layer is washed with dichloromethane (250 ml). Dichloromethane (1

-35-

L) is added to the acidic aqueous phase and the pH is adjusted to 12-13 with 5N aqueous sodium hydroxide. The phases are separated and the organic phase is dried over sodium sulfate. After filtration the solution is
5 concentrated to give 53 g of the title amine (88%).

Amine 6

4-(2-Amino-2-methylpropyl)phenol (3.00 g, 18.2
10 mmol), methyl 6-chloronicotinate (3.27 g, 19.1 mmol),
powdered potassium carbonate (3.76 g, 27.2 mmol, 300 mesh),
N,N-dimethylacetamide (60 ml), and toluene (15 ml) are
combined and heated to reflux. The water formed during the
reaction is removed azeotropically via a Dean-Stark trap.
15 After about 2 hours, the internal temperature reached 154°C
and the reaction is complete. The slurry is cooled to 30°C
and filtered. The filter cake is washed with N,N-
dimethylacetamide and the combined organic fractions
concentrated by rotary evaporation at 75°C. The resulting
20 oil is dissolved in ethyl acetate (50 ml), and washed with
water (30 ml). The phases are separated and the aqueous
phase is extracted with ethyl acetate (20 ml) after some
saturated aqueous sodium chloride solution (10 ml) is added
to facilitate phase separation. The combined organic
25 fractions are washed with water (2 x 30 ml) and saturated
aqueous sodium chloride (30 ml) and then dried over sodium
sulfate. After filtration the solution is concentrated to
give 4.60 g (80%) of the title amine.

30

Amine 7

4-(2-Amino-2-methylpropyl)phenol acetic acid salt
(45.1 g, 0.200 mol), powdered potassium carbonate (58.1 g,
0.420 mol, 300 mesh), and isooctane (140 mL) are added

-36-

sequentially to N,N-dimethylacetamide (500 mL) and the resulting mixture is heated to reflux under a Dean-Stark trap until water production almost ceased (about 2 hours, 121°C internal temperature). A solution of ethyl 6-chloronicotinate (39.0 g, 0.210 mol) in N,N-dimethylacetamide (50 mL) is added over 15 minutes and the reaction heated at reflux, continuing to remove any water or alcohol as formed, until the reaction is complete by HPLC (about 2 hours). The slurry is cooled to room temperature and filtered through a one quarter inch pad of celite and the cake is rinsed with methyl tert-butyl ether (2 x 75 mL). The filtrate is then concentrated by rotary evaporation to a net weight of about 134 g. The resulting oil is dissolved in methyl tert-butyl ether (315 mL) and washed with water (315 mL). The product is extracted from the organic phase with aqueous hydrochloric acid (1N,, 220 mL) and water (140 mL). Ethyl acetate (315 mL) is added to the aqueous phase and the mixture basified with a solution of sodium carbonate (1.2 equivalents in 135 mL water) with stirring. Caution: foaming may occur. The phases are separated and the organic phase is washed with 10% w/w aqueous solution of sodium chloride (135 mL). The solvent is removed by rotary evaporation, the oil is dissolved in methanol (135 mL), and the solvent concentrated again. The resulting solid is dried overnight at room temperature to afford 61.0 g of the title amine (94% yield) which is used without further purification.

Amine 11

2-Cyano-3-chloropyridine (Bremner, et al., *Syn. Comm.*, 27:1535, 1997; Kaneda, et al., *Chem. Pharm. Bull.*, 33:565, 1985) is coupled to 4-(2-amino-2-methylpropyl)phenol

-37-

to prepare the title amine by a procedure substantially similar to that described above for Amine 4.

Amine 23

5

Potassium tert-butoxide (58.6 ml, 58.6 mmol, 1M in tetrahydrofuran) is added to a solution of 3,4-dichlorothiophenol (10.0 g, 55.8 mmol) in tetrahydrofuran (300 ml) at 0°C and the solution stirred for 30 minutes.

10 Methyl iodide (8.32 g, 58.6 mmol) is added dropwise and the resulting slurry is stirred for 16 hours. The solvents are removed *in vacuo* and the residue is dissolved in 150 ml each of methyl-t-butyl ether and 1M NaHSO₄. The phases are separated and the organic layer is washed with 150 ml each
15 of water and saturated aqueous sodium chloride. The organic layer is dried over sodium sulfate, filtered and concentrated *in vacuo* to give 9.67 g of 3,4-dichlorophenyl methylsulfide (90%).

The sulfide from above is converted to the
20 corresponding sulfone as described below for Amine 37. The 3,4 dichloromethyl sulfone is coupled to 4-(2-amino-2-methylpropyl)phenol to prepare the title amine by a procedure substantially similar to that described above for Amine 4.

25

Amine 30

To a 1 gallon autoclave is added 2,5-dichloropyridine (123 g, 830 mmol), palladium II acetate
30 (5.6 g, 24.9 mmol), 1,3-bis(diphenylphosphine)propane (20.5 g, 49.8 mmol), 1,1,1,3,3,3,-hexamethyldisilazane (700 ml), acetonitrile (1180 ml) and dimethylformamide (295 ml). The autoclave is pressurized to 70 psi with carbon monoxide and heated to 80°C for 16 hours. The reaction mixture is

-38-

filtered and washed with acetonitrile. The mixture is concentrated *in vacuo* to 590 g and 1L of water is added. The resulting slurry is cooled to 0°C and filtered to give 102.6 g (79%) of 2-carboxamido-5-chloropyridine which is used without further purification.

2-Carboxamido-5-chloropyridine is coupled to 4-(2-amino-2-methylpropyl)phenol to prepare the title amine by a procedure substantially similar to that described above for Amine 4.

Amine 33

4-Chloro-1,3-benzenedithiol (3.06 g, 17.3 mmol) is added to a stirred mixture of potassium carbonate (5.20 g, 37.7 mmol) in 10 ml of anhydrous dimethylformamide at room temperature under a nitrogen atmosphere. After five minutes, iodomethane (7.20 g, 50.7 mmol) is added in a dropwise fashion to give an exothermic reaction. The reaction mixture is stirred at room temperature over night. The mixture is then diluted with 50 ml deionized water and extracted with hexanes (3 x 30 ml). The combined hexane extracts are washed with brine and dried over anhydrous magnesium sulfate. Filtration and evaporation of the solvent gave an oil. This oil is dissolved in 50 ml of glacial acetic acid with stirring and chilled to 0°C in an ice/water bath. Hydrogen peroxide (30%, 8 ml) is added slowly and the mixture stirred at 0°C for 15 minutes. The ice bath is removed and the mixture allowed to warm to room temperature for one hour. It is then heated under reflux over night. The reaction mixture is concentrated until a precipitate formed. The mixture is then diluted with 50 ml methylene chloride and 50 ml water, neutralized with saturated sodium bicarbonate solution, and the organic layer collected. The aqueous layer is again extracted with

-39-

methylene chloride (2 x 50ml) and the combined organic layers washed with brine and dried over anhydrous magnesium sulfate. After filtration and concentration, 4.29 g of methyl(4-chloro-3-methylsulfonylphenyl)sulfone is obtained
5 (92%). Methyl(4-chloro-3-methylsulfonylphenyl)sulfone is converted to the title amine by a procedure substantially analogous to that described for Amine 9.

Amine 35

10

To a solution of n-butyl lithium (0.544 mol) in tetrahydrofuran (700 ml) at -78°C is added a solution of 3,4-difluorobromobenzene (100 g, 0.518 mol) in 200 ml of tetrahydrofuran. After 10 minutes, a solution of dimethyl
15 disulfide in 100 ml of tetrahydrofuran is added and the resulting reaction mixture is warmed to ambient temperature over 60 minutes. The reaction is concentrated *in vacuo* and the resulting oil is partitioned between 750 ml methyl-t-butyl ether and 300 ml water. The phases are separated and
20 the organic layer is washed with 300 ml of saturated aqueous sodium chloride, dried over magnesium sulfate, filtered, and concentrated *in vacuo*. The resulting oil is purified by vacuum distillation to provide 43.08 g of 3,4-difluorophenyl methylsulfide.

25

Metachloroperbenzoic acid (60.4 mmol) is added portionwise to a solution of the sulfide (43 g, 26.8 mmol) in 1L of dichloromethane at 0°C. After 15 minutes, the reaction mixture is warmed to ambient temperature and stirred for 1.25 hours. The solids are removed by
30 filtration and the resulting solution washed with 750 ml of 1M sodium bisulfite, 2L sodium bicarbonate, 1L water, and 750 ml saturated aqueous sodium chloride. The organic layer is dried over magnesium sulfate, filtered and concentrated *in vacuo* to give 45.17 g (88%) of 2,4-difluorophenyl methyl

-40-

sulfone. The sulfone is converted to the title amine by a procedure substantially analogous to that described for Amine 9.

5 Amines 5, 12-22, 24, 27, 29, 31, 34 and 36

Amines 5, 12-22, 24, 27, 29, 31, 34 and 36 are prepared by procedures substantially similar to that described for Amine 4.

10

Examples

Representative Procedure 3: Amination of Epoxide

15 A vial is charged with a solution of single amine of formula III (0.2M in ethanol or t-butanol, 90 micromolar) and a solution of a single epoxide of formula II (0.2M in dimethylsulfoxide, 80 micromolar). The vial is sealed and heated to 80°C for 24-48 hours. The solution is cooled to
20 room temperature, diluted with methanol, and passed over a cation exchange column, eluting the basic material with 1N methanolic ammonia.

Representative Procedure 4: Amination of Epoxide

25

A stirred mixture of an epoxide of formula II (1 equivalent) and an amine of formula III (1-2 equivalents) in ethanol, methanol, n-butanol or t-butanol is heated at 70-80°C for 2-72 hours. The solvent is evaporated to dryness
30 to give a crude oil that is optionally diluted with methanol or ethanol and passed over a cation exchange column (eluting the free base product with 1N methanolic ammonia) before further purification.

-41-

The final products prepared via Representative Procedure 3 or 4 may be further purified by flash or radial chromatography. Typical chromatography conditions include:

- a) using a variable mixture of 25:5:1
- 5 chloroform/methanol/ammonium hydroxide and 9:1 chloroform/methanol; b) a variable mixture of 90:10:1 CH_2Cl_2 /ethanolic NH_3 gradient; c) dichloromethane/6-12% methanol, 0.15-0.35M ammonia in dichloromethane gradient; d) methylene chloride with a step gradient to 2-8% methanol; e)
- 10 chloroform/2.0M ammonia in methanol, from 0-10% to 6-20% gradient elution or f) isocratic 6-8% 2M ammonia in methanol: 92-94% dichloromethane.

Alternatively, the final products may be purified on C18 bonded silica gel using either mass guided or UV

15 guided reverse phase liquid chromatography (acetonitrile/water with 0.01% hydrochloric acid or 0.1% trifluoroacetic acid). When purification of a compound of the present invention results in production of a free base, the free base thus prepared may be salified, e.g., by

20 dissolution of the free base in CH_2Cl_2 or diethylether, adding 1M ethanolic HCl or a solution of HCl in diethylether, and evaporating the volatiles, or as described in more detail below.

For example, a hydrochloride salt may be prepared

25 by dissolving the free base in dichloromethane, diethylether, or a mixture of ethyl acetate and methanol and adding 1M ethanolic HCl, a solution of HCl in diethylether, or 0.5M ammonium chloride. The resulting mixture is allowed to stir for a short time, e.g., for five minutes, before

30 evaporating the volatiles and optionally triturating in diethyl ether to give the hydrochloride salt.

The oxalate salts may be prepared by dissolving the free base in a small amount of ethyl acetate, optionally adding methanol for solubility. The resulting solution is

-42-

treated with 1 equivalent of a 0.5M solution of oxalic acid in ethyl acetate. The reaction mixture is either concentrated *in vacuo* or centrifuged, separated, and the solids are dried, to give the oxalate salt.

5 To prepare a succinate salt, the free base may be dissolved in a small amount of ethyl acetate or methanol and then treated with 1 equivalent of succinic acid in methanol. The resulting slurry is dissolved in the minimum amount of methanol then concentrated *in vacuo* to give the succinate
10 salt.

For products synthesized from Amine 38, after amination of the epoxide, the crude ethanolamine product is further treated with 1M tetrabutyl ammonium fluoride (to remove the t-butyldimethylsilyl protecting group) in
15 tetrahydrofuran and stirred at room temperature in tetrahydrofuran overnight. The product is then isolated as described above.

The table below sets out representative combinations of Amines and Epoxides that are reacted as
20 described above in Representative Procedure 3 or 4. Preparation of desired product is confirmed via mass spectral analysis (MSA). Emax \pm Standard Error Mean (SEM) data, discussed in the "Demonstration of Function" section below, is also included for said compounds where available.
25 The Emax values represent the average of at least 3 runs except as otherwise indicated.

Table 3

E.g.	Epoxide	Amine	MSA	Isolated Form	Emax (%) \pm SEM
1	1	8	504.9	Free Base	72.6 \pm 5.4
2	2	1	476.3	Trifluoroacetate	58.3 \pm 5.7
3	2	1	476.3	Hydrochloride	Not Tested
4	2	1	476.3	Free Base	Not Tested
5	2	2	501.3	Hydrochloride	61.0 \pm 4.5

Table 3 (continued)

6	2	3	519.0	Free Base	67.9 ± 2.0
7	2	4	501.3	Hydrochloride	64.0 ± 3.6
8	2	5	519.3	Hydrochloride	67.9 ± 1.4
9	2	7	548.3	Trifluoroacetate	Not Tested
10	2	8	518.3	Trifluoroacetate	68.3 ± 2.3
11	2	9	553.3	Hydrochloride	59.3 ± 2.6
12	2	15	578.2	Trifluoroacetate	60.6 ± 8.6
13	2	16	534.2	Trifluoroacetate	51.7 ± 4.3
14	2	17	552.3	Trifluoroacetate	57.2 ± 3.1
15	2	18	510.3	Trifluoroacetate	65.1 ± 2.6
16	2	19	577.3	Trifluoroacetate	47.4 ± 2.7
17	2	20	544.2	Trifluoroacetate	52.7 ± 4.5
18	2	21	510.2	Trifluoroacetate	56.2 ± 1.8
19	2	22	510.2	Trifluoroacetate	61.7 ± 1.2
20	2	28	519.3	Trifluoroacetate	69.8 ± 5.4
21	2	30	519.2	Trifluoroacetate	67.1 ± 4.5
22	2	31	511.2	Trifluoroacetate	72.0 ± 1.6
23	2	32	518.2	Free Base	80.4 ± 7.0
24	2	34	553.2	Free Base	66.8 ± 6.4
25	2	35	571.2	Trifluoroacetate	54.5 ± 4.9
26	2	36	572.2	Trifluoroacetate	55.3 ± 5.2
27	3	3	553.3	Hydrochloride	41.7 ± 8.4
28	3	5	533.4	Hydrochloride	49.7 ± 5.1
29	4	3	547.3	Hydrochloride	49.4 ± 5.7
30	5	3	581.3	Hydrochloride	15.7 ± 0.6
31	6	3	517.3	Hydrochloride	76.8 ± 3.3
32	6	5	517.2	Hydrochloride	73.9 ± 1.3
33	7	3	531.2	Free Base	71.2 ± 3.9
34	8	1	502.3	Trifluoroacetate	56.3 ± 4.6
35	8	2	527.3	Free Base	54.6 ± 4.7
36	8	2	527.3	Hydrochloride	49.9 ± 1.7
37	8	3	545.3	Free Base	39.3 ± n = 1
38	8	3	545.3	Hydrochloride	66.3 ± 2.9
39	8	4	527.3	Free Base	67.6 ± 3.1
40	8	4	527.3	Hydrochloride	70.6 ± 4.5
41	8	5	545.4	Free Base	Not Tested
42	8	5	545.3	Hydrochloride	67.3 ± 9.4
43	8	6	560.4	Free Base	63.6 ± 4.0
44	8	7	574.1	Free Base	64.3 ± 4.1
45	8	9	579.3	Free Base	59.1 ± 4.4
46	8	9	579.2	Hydrochloride	48.8 ± 4.2
47	8	10	593.3	Free Base	51.4 ± 2.8
48	8	11	527.3	Trifluoroacetate	51.6 ± 1.8
49	8	12	503.2	Trifluoroacetate	60.6 ± 2.9
50	8	13	503.3	Trifluoroacetate	47.5 ± 3.4
51	8	14	606.2	Trifluoroacetate	42.5 ± 3.1
52	8	17	578.2	Trifluoroacetate	47.1 ± 4.7

-44-

Table 3 (continued)

53	8	20	570.2	Trifluoroacetate	51.3 ± 3.8
54	8	21	536.2	Trifluoroacetate	47.6 ± 3.0
55	8	22	536.2	Trifluoroacetate	44.9 ± 6.0
56	8	23	613.2	Trifluoroacetate	38.4 ± 3.3
57	8	24	591.3	Trifluoroacetate	36.8 ± 2.7
58	8	25	544.3	Trifluoroacetate	35.7 ± 0.8
59	8	26	562.3	Trifluoroacetate	44.3 ± 5.3
60	8	27	570.3	Trifluoroacetate	50.1 ± 2.7
61	8	28	545.3	Trifluoroacetate	64.6 ± 1.2
62	8	29	502.3	Trifluoroacetate	52.6 ± 3.8
63	8	30	545.3	Trifluoroacetate	69.9 ± 8.1
64	8	30	545.2	Free Base	Not Tested
65	8	31	537.2	Trifluoroacetate	54.7 ± 5.1
66	8	32	544.3	Free Base	43.5 ± 3.9
67	8	33	657.2	Free Base	58.7 ± 5.2
68	8	34	579.2	Free Base	66.0 ± 7.4
69	8	35	597.2	Trifluoroacetate	48.6 ± 5.1
70	8	36	598.2	Trifluoroacetate	52.5 ± 4.5
71	8	37	425.2	Free Base	46.5 ± 3.5
72	8	38	544.3	Free Base	74.7 ± 3.0
73	9	3	559.3	Hydrochloride	41.0 ± 5.7
74	9	5	559.3	Hydrochloride	50.8 ± 9.8
75	7	5	531.2	Hydrochloride	82.0 ± 2.5
76	7	30	531.2	Hydrochloride	71.4 ± 10.6

Alternate Preparation of Example 37

5 A 1 gallon autoclave is charged with the compound
of Example 45 (105 g, 183 mmol) and methanol (1.2 L). The
tank is pressurized with ammonia gas to 50 psig and the
contents are heated to 40°C. After 66 hours the tank is
cooled to 20°C, and the solution is filtered through celite
10 and glass paper. The tank is rinsed with a total of 1.5 L
of methanol and the combined filtrates are concentrated to a
foam (105.41 g). MS (ES+) m/z 545.

Alternate Preparation of Example 38

15

The compound of Example 37 (160.20 g, 294.1 mmol)
is dissolved in methanol (1.6 L). Concentrated hydrochloric
acid (24.5 mL) is added and the resulting solution is heated

-45-

to 60-65°C. Methyl-t-butylether is then added (1.2 L) over the course of 2.25 hours while maintaining reflux. When the addition is complete the mixture is allowed to cool to 21°C, at which point it is further cooled to < 2°C in an ice bath.

- 5 The mixture is stirred for 1 hour and is then filtered. The solids are dried in a 50°C vacuum oven to provide 133.61 g (78%) of crystalline solid. MS (ES+) m/z 545.

Alternative Preparation of Example 44

10

- To a 3 neck, 3 L flask, Epoxide 8 (122 g, 469 mmol, 1 equivalent), Amine 7 (250 g, 797 mmol, 1.7 equivalents), and ethanol denatured with toluene (1.5 L, 12 vol.) is added. The reaction is heated to reflux and
15 allowed to stir at reflux for 25 hours. The reaction is then concentrated in vacuum to oil. The oil is taken up in ethyl acetate (10 volumes, 2x) and stripped down to a glassy oil. The oil is then dissolved in ethyl acetate (10 volumes) and water (5 volumes). The two layers are
20 separated and the resulting organic layer is washed with water (5 volumes based on product). The resulting organic layer is then washed with 1N aqueous hydrochloric acid / water (0.25 eq of acid based on excess amine). The organic layer is then checked, via HPLC, for any excess amine. If
25 amine is detected the organic layer is washed with 1N aqueous hydrochloric acid/water (increments of 0.1 eq of acid) until all the amine is completely removed. The organic layer is washed with 1N aqueous sodium hydroxide (1 volume) and saturated sodium chloride (5 vol.), dried over
30 magnesium sulfate, filtered, and stripped to a foam. The foam is chromatographed using 3 kg silica gel. Eluent consisted of 3A ethanol:tetrahydrofuran:heptane:dichloromethane (first 8 fractions, 3:6:31:60, next 5

-46-

4:8:28:60, last 5 fractions 7.5:12.5:20:60). MS (ES+) m/z 574.1.

Demonstration of Function

5 The genes encoding the human β_1 -adrenergic receptor (Frielle et al., *Proc. Natl. Acad. Sci.*, 84:7920-7924, 1987), the human β_2 -adrenergic receptor (Kobika et al., *Proc. Natl. Acad. Sci.*, 84:46-50, 1987, Emorine et al., *Proc. Natl. Acad. Sci.*, 84:6995-6999, 1987) and the human β_3 10 adrenergic receptor (Granneman et al., *Molecular Pharmacology*, 44(2):264-70, 1993) are individually subcloned into a phd expression vector (Grinnell et al., *Bio/Technology*, 5:1189-1192, 1987) and transfected into the DXB-11 Chinese hamster ovary (CHO) cell line by calcium 15 phosphate precipitation methodology. The stably transfected cells are grown to 95% confluency in 95% Dulbecco's modified Eagles Medium (DMEM), 5% fetal bovine serum and 0.01% proline. Media is removed and the cells are washed with phosphate buffered (pH 7.4) saline (without magnesium and 20 calcium). Cells are then lifted using an enzyme free cell dissociation solution (Specialty Media, Lavallette, New Jersey) and pelleted by centrifugation.

 Cells from each of the above cell lines are resuspended and added (20,000/well) to a 96-well plate. 25 Cells are incubated at 37°C with representative compounds of the invention for 20 minutes in buffer (Hank's balanced salt solution, 10 mM HEPES, 0.1% BSA, 1 mM L-ascorbic acid, 0.2% dimethyl sulfoxide, 1 mM 3-isobutyl-1-methylxanthine, pH 7.4). After halting the incubation with quench buffer (50 30 mM Na Acetate, 0.25% Triton X-100, pH 5.8), the c-AMP level is quantified by scintillation proximity assay (SPA) using a modification of the commercially available c-AMP kit

-47-

(Amersham, Arlington Heights, IL) with rabbit anti-cAMP antibody (ICN Biomedicals, Aurora, Ohio) for the kit.

Sigmoidal dose response curves, from the whole cell receptor coupled c-AMP assay are fit to a four
5 parameter logistic equation using non linear regression:
 $y = (a-d) / (1 + (Dose/c)^b) + d$ where a and d are responses at zero and maximal dose, b is the slope factor and c is the EC₅₀ as previously described (DeLean et al., *Am. J. Physiol.*, 235, E97-E102, 1978). EC₅₀ is assessed as the concentration
10 producing 50% of the maximum response to each agonist.

Isoproterenol is accepted in the art as a non-selective β_3 agonist and is widely used as a comparator in evaluating the activity of compounds. See *Trends in Pharm. Sci.*, 15:3, 1994. The % intrinsic activity (E_{max}) of
15 representative compounds of the invention is assessed relative to isoproterenol by the compound's maximal response divided by the isoproterenol maximal response times 100.

In vitro Rat Atrial Tachycardia

20 Male rats (250-350 g) (Harlan Sprague Dawley, Indianapolis, Indiana, USA) are killed by cervical dislocation. Hearts are removed and the left and right atria are dissected and mounted with thread in tissue baths containing 10 mls of modified Krebs' solution. Initial
25 resting tension is 1.5-2.0 g at the outset of the experiment (Naunyn-Schmied Arch. Pharmacol., 320:145, 1982). Tissues are allowed to equilibrate approximately 30 minutes with vigorous oxygenation before exposure to a compound of the invention.

30 To evaluate the ability of test compounds to increase heart rate, representative compounds of the present invention are added cumulatively once the atrial rate reached a steady state from the previous addition. Compound

-48-

addition is continued until no further increase in atrial rate occurred or until a concentration of 10^{-4} M is reached. The increase in beats per minute (bpm) is measured for each concentration of test compound by means of a BioPac System
5 (Br. J. of Pharmacol., 126:1018-1024, 1999).

Relative to a compound of formula I where R^1 , R^{1a} , R^2 , R^3 and R^4 are hydrogen, R^5 and R^6 are both methyl, X^2 is O, and R^7 is 5-carboxamido-pyrid-2-yl (not claimed, generically disclosed in U.S. Patent No. 5,786,356),
10 corresponding compounds of the present invention exhibit a reduction in atrial tachycardia.

Utilities

As agonists of the β_3 recptor, a compound of the
15 present invention is useful in treating conditions in human and non-human animals in which the β_3 receptor has been demonstrated to play a role. The diseases, disorders or conditions for which compounds of the present invention are useful in treating or preventing include, but are not
20 limited to, (1) diabetes mellitus, (2) hyperglycemia, (3) obesity, (4) hyperlipidemia, (5) hypertriglyceridemia, (6) hypercholesterolemia, (7) atherosclerosis of coronary, cerebrovascular and peripheral arteries, (8) gastrointestinal disorders including peptid ulcer,
25 esophagitis, gastritis and duodenitis, (including that induced by H. pylori), intestinal ulcerations (including inflammatory bowel disease, ulcerative colitis, Crohn's disease and proctitis) and gastrointestinal ulcerations, (9) neurogenic inflammation of airways, including cough, asthma,
30 (10) depression, (11) prostate diseases such as benign prostate hyperplasia, (12) irritable bowel syndrome and other disorders needing decreased gut motility, (13) diabetic retinopathy, (14) neuropathic bladder dysfunction,

-49-

(15) elevated intraocular pressure and glaucoma and (16) non-specific diarrhea dumping syndrome.

In treating non-human, non-companion animals, the compounds of the present invention are useful for increasing weight gain and/or improving the feed utilization efficiency and/or increasing lean body mass and/or decreasing birth mortality rate and increasing post/natal survival rate.

Formulation

10 The compound of formula I is preferably formulated in a unit dosage form prior to administration. Therefore, yet another embodiment of the present invention is a pharmaceutical formulation comprising a compound of formula I and a pharmaceutical carrier.

15 The present pharmaceutical formulations are prepared by known procedures using well-known and readily available ingredients. In making the formulations of the present invention, the active ingredient (formula I compound) will usually be mixed with a carrier, or diluted by a carrier, or enclosed within a carrier which may be in the form of a capsule, sachet, paper or other container. When the carrier serves as a diluent, it may be a solid, semisolid or liquid material which acts as a vehicle, excipient or medium for the active ingredient. Thus, the compositions can be in the form of tablets, pills, powders, lozenges, sachets, cachets, elixirs, suspensions, emulsions, solutions, syrups, aerosol (as a solid or in a liquid medium), soft and hard gelatin capsules, suppositories, sterile injectable solutions and sterile packaged powders.

30 Some examples of suitable carriers, excipients, and diluents include lactose, dextrose, sucrose, sorbitol, mannitol, starches, gum acacia, calcium phosphate, alginates, tragacanth, gelatin, calcium silicate, microcrystalline cellulose, polyvinylpyrrolidone, cellulose,

-50-

- water syrup, methyl cellulose, methyl and propylhydroxybenzoates, talc, magnesium stearate and mineral oil. The formulations can additionally include lubricating agents, wetting agents, emulsifying and suspending agents, preserving agents, sweetening agents or flavoring agents. The compositions of the invention may be formulated so as to provide quick, sustained or delayed release of the active ingredient after administration to the patient.

10

Formulation ExamplesFormulation 1: Tablets

Ingredient	Quantity (mg/tablet)
Active Ingredient	5 - 500
Cellulose, microcrystalline	200 - 650
Silicon dioxide, fumed	10 - 650
Stearate acid	5 - 15

- 15 The components are blended and compressed to form tablets.

Formulation 2: Suspensions

Ingredient	Quantity (mg/5 ml)
Active Ingredient	5 - 500 mg
Sodium carboxymethyl cellulose	50 mg
Syrup	1.25 mg
Benzoic acid solution	0.10 ml
Flavor	q.v.
Color	q.v.
Purified water to	5 ml

- 20 The medicament is passed through a No. 45 mesh U.S. sieve and mixed with the sodium carboxymethyl cellulose and syrup

-51-

to form a smooth paste. The benzoic acid solution, flavor, and color are diluted with some of the water and added, with stirring. Sufficient water is then added to produce the required volume.

5

Formulation 3: Intravenous Solution

Ingredient	Quantity
Active Ingredient	25 mg
Isotonic saline	1,000 ml

The solution of the above ingredients is intravenously administered to a patient at a rate of about 1 ml per minute.

Dose

The specific dose administered is determined by the particular circumstances surrounding each situation. These circumstances include, the route of administration, the prior medical history of the recipient, the pathological condition or symptom being treated, the severity of the condition/symptom being treated, and the age and sex of the recipient. However, it will be understood that the therapeutic dosage administered will be determined by the physician in the light of the relevant circumstances.

Generally, an effective minimum daily dose of a compound of formula I is about 5, 10, 15, or 20 mg. Typically, an effective maximum dose is about 500, 100, 60, 50, or 40 mg. Most typically, the dose ranges between 15 mg and 60 mg. The exact dose may be determined, in accordance with the standard practice in the medical arts of "dose titrating" the recipient; that is, initially administering a low dose of the compound, and gradually increasing the dose until the desired therapeutic effect is observed.

-52-

Route of Administration

The compounds can be administered by a variety of routes including the oral, rectal, transdermal,
5 subcutaneous, topical, intravenous, intramuscular or intranasal routes.

Combination Therapy

10 A compound of formula I may be used in combination with other drugs that are used in the treatment/prevention/suppression or amelioration of the diseases or conditions for which compounds of formula I are useful. Such other drug(s) may be administered, by a route
15 and in an amount commonly used therefor, contemporaneously or sequentially with a compound of formula I. When a compound of formula I is used contemporaneously with one or more other drugs, a pharmaceutical unit dosage form containing such other drugs in addition to the compound of
20 formula I is preferred. Accordingly, the pharmaceutical compositions of the present invention include those that also contain one or more other active ingredients, in addition to a compound of formula I. Examples of other active ingredients that may be combined with a compound of
25 formula I, either administered separately or in the same pharmaceutical compositions, include, but are not limited to:

(a) insulin sensitizers including (i) PPAR γ agonists such
30 as the glitazones (e.g. troglitazone, pioglitazone, englitazone, MCC-555, BRL49653 and the like), and compounds disclosed in WO97/27857, 97/28115, 97/28137 and 97/27847; (ii) biguanides such as metformin and phenformin;

-53-

- (b) insulin or insulin mimetics;
- (c) sulfonylureas such as tolbutamide and glipizide;
- (d) alpha-glucosidase inhibitors (such as acarbose);
- (e) cholesterol lowering agents such as
 - 5 i. HMG-CoA reductase inhibitors (lovastatin, simvastatin and pravastatin, fluvastatin, atorvastatin, and other statins),
 - ii. sequestrants (cholestyramine, colestipol and a dialkylaminoalkyl derivatives of a cross-linked
10 dextran),
 - iii. nicotiny alcohol nicotinic acid or a salt thereof,
 - iv. proliferator-activator receptor agonists such as fenofibric acid derivatives (gemfibrozil, clofibrat, fenofibrate and benzafibrate),
15 v. inhibitors of cholesterol absorption for example beta-sitosterol and (acyl CoA:cholesterol acyltransferase) inhibitors for example melinamide,
 - 20 vi. probucol,
 - vii. vitamin E, and
 - viii. thyromimetics;
- (f) PPAR δ agonists such as those disclosed in
25 WO97/28149;
- (g) antiobesity compounds such as fenfluramine, dexfenfluramine, phentermine, sibutramine, orlistat, and other β_3 adrenergic receptor agonists;
- (h) feeding behavior modifying agents such as
30 neuropeptide Y antagonists (e.g. neuropeptide Y5) such as those disclosed in WO 97/19682, WO 97/20820, WO 97/20821, WO 97/20822 and WO 97/20823;


-54-

- (i) PPAR α agonists such as described in WO 97/36579 by Glaxo;
- (j) PPAR γ antagonists as described in WO97/10813; and
- (k) serotonin reuptake inhibitors such as fluoxetine and
5 sertraline.

-55-

WE CLAIM:

1. A compound of formula I:



I;

5 wherein:

R^1 is H, CN, halo, C_1 - C_6 alkyl, C_1 - C_4 haloalkyl, CO_2R^8 , $CONHR^8$, $NHCO_2R^8$, NHR^8 , OR^8 , SR^8 , SOR^8 , SO_2R^8 or SO_2NHR^8 ;

R^{1a} is H, halo or C_1 - C_6 alkyl;

10 R^2 is H, C_1 - C_6 alkyl or benzyl;

R^3 is C_1 - C_6 alkyl or benzyl;


or R^2 and R^3 combine with the carbon to which each are attached to form a C_3 - C_7 carbocyclic ring; provided that if R^3 is C_2 - C_6 alkyl or benzyl, then R^2 must be hydrogen;

15 R^4 is H or C_1 - C_6 alkyl;

R^5 and R^6 are independently H or C_1 - C_6 alkyl; or

R^5 and R^6 combine with the carbon to which each are attached to form a C_3 - C_6 carbocyclic ring;

20 or R^6 combines with X^1 , the carbon to which both are attached, and the phenyl group to which X^1 is attached to form a moiety selected from the group consisting of:



wherein:

-56-

m and n are independently 0, 1, 2, or 3
provided that the sum of n + q is ≤ 5 and
that R⁵ is H;

R⁷ is hydrogen, optionally substituted phenyl or
5 optionally substituted heterocycle;

R⁸ is H or C₁-C₆ alkyl;

X is OCH₂, SCH₂ or a bond; and

X¹ is a bond or a C₁-C₅ divalent hydrocarbon
moiety; and

10 X² is O, S, NH, NHSO₂, SO₂NH, CH₂ or a bond;
or a pharmaceutical salt thereof.

2. The compound of Claim 1 wherein:

R¹ is H, CN, halo, C₁-C₄ alkyl, C₁-C₄ haloalkyl,
15 CO₂R⁸, CONHR⁸, NHCOR⁸, NHR⁸, OR⁸, SR⁸, SOR⁸, SO₂R⁸ or
SO₂NHR⁸;

R^{1a} is H, halo or C₁-C₄ alkyl;

R⁴ is H or C₁-C₄ alkyl;

R⁵ and R⁶ are independently H or C₁-C₄ alkyl; or
20 R⁵ and R⁶ combine with the carbon to which each are
attached;

or R⁶ combines with X¹, the carbon to which both
are attached, and the phenyl group to which X¹ is attached;

R⁷ is hydrogen, phenyl or heterocycle wherein said
25 phenyl or heterocycle is optionally substituted one to three
times independently with hydroxy, oxo, nitro, phenyl,
benzyl, C₁-C₄ alkoxy, COR⁸, NHCO(C₁-C₄ alkyl), NHCO(phenyl),
NHCO(benzyl), OCO(C₁-C₄ alkyl), OCO₂R⁸ and OCONR⁸R⁸; and

-57-

R^8 is H or C_1 - C_4 alkyl; or a pharmaceutical salt thereof.

3. The compound of Claim 1 or 2 wherein:

5

R^1 is H;

R^{1a} is H;

R^2 and R^3 are both methyl or combine with the carbon to which each are attached to form a pentacyclic ring;

10

R^4 is H;

R^5 and R^6 are independently H or methyl;

R^7 is phenyl, pyridyl, pyridazinyl or pyrimidinyl wherein said R^7 moieties are optionally substituted once or twice with chloro, cyano, $CONH_2$ or CO_2CH_3 ;

15


X is OCH_2 and is connected to the indole ring system at the 4-position of said system;

X^1 is CH_2 ; and

X^2 is O; or a pharmaceutical salt thereof.

20

4. The compound of any one of Claims 1-3 of the formula:




wherein R^7 is phenyl or pyridyl and said R^7 moieties are substituted once with cyano or $CONH_2$; or a pharmaceutical salt thereof.

25

-58-

5. The compound of any one of Claims 1-4 which is selected from the group consisting of:



or a pharmaceutical salt thereof.

5

6. The compound of any one of Claims 1-5 which is the hydrochloride salt.

10

7. A pharmaceutical formulation comprising a compound of any one of Claims 1-6 and a pharmaceutical carrier.

15

8. A method of treating Type II Diabetes comprising administering to a patient in need thereof a compound of any one of Claims 1-6.

20


9. A method of treating obesity comprising administering to a patient in need thereof a compound of any one of Claims 1-6.

-59-

10. A method of agonizing the β_3 receptor comprising administering to a patient in need thereof a compound of any one of Claims 1-6.

5 11. A compound of any one of Claims 1-6 for use in treating Type II diabetes, obesity or for use in agonizing the β_3 receptor.

12. A compound of formula II:



10

wherein:

R^1 is H, CN, halo, C_1 - C_6 alkyl, C_1 - C_4 haloalkyl, CO_2R^8 , $CONHR^8$, $NHCO_2R^8$, NHR^8 , OR^8 , SR^8 , SOR^8 , SO_2R^8 or SO_2NHR^8 ;


15 R^2 is H, C_1 - C_6 alkyl or benzyl;

R^3 is C_1 - C_6 alkyl or benzyl;

R^8 is H or C_1 - C_6 alkyl; and

X is OCH_2 , SCH_2 or a bond; or a salt thereof.


20 13. The compound of Claim 12 of the formula:



or a salt thereof.


-60-

14. A process for preparing a compound of any one of Claims 1-6 which comprises reacting a compound of formula II:



II;

with a compound of formula III:



III;

in the presence of a suitable solvent.